

pathognomonic of leukemia but is now recognized as a sign of any chronic illness with bone growth retardation.

Growth lines can often be correlated with a definitely marked illness but there is no good correlation with the severity of the illness, for often severe illnesses leave no such marks. Conversely, many children will show growth lines during a period in which no cessation of growth or illness can be documented.

M. B. OZONOFF, M.D.

REFERENCES

- Garn SM, Silverman FN, Hertzog KP, et al: Lines and bands of increased density. Their implication to growth and development. *Med Radiogr Photogr* 44:58-89, 1968
- Park EA: The imprinting of nutritional disturbances on the growing bone. *Pediatrics* 33(Supplement):815-862, 1964
- Wells C: *Bones, bodies and disease*. London, Thames & Hudson, 1964

Congenital Bone Lesions Following Fetal Viremia

The longitudinal striations seen in the metaphyseal portions of some long bones in rubella syndrome are now thought to be due to damage to the fetal mesoderm with failure of subsequent maturation to osteoblasts. Consequently, columns of defective bone are formed which appear roentgenographically at birth as longitudinal, vertical, lucent striations. They are most common in the humeri and femurs. They usually disappear through remodeling in a few months.

These lesions were originally reported only with rubella syndrome, but recently reports have documented identical lesions in cytomegalic inclusion disease.

M. B. OZONOFF, M.D.

REFERENCES

- Rudolph AJ, Singleton EB, Rosenberg HS, et al: Osseous manifestations of the congenital rubella syndrome. *Amer J Dis Child* 110:428-443, 1965
- Graham CB, Thal A, Wassum CS: Rubella-like bone changes in congenital cytomegalic inclusion disease. *Radiology* 94:39-43, 1970
- Reed GP Jr: Rubella bone lesions. *J Pediatr* 74:208-213, 1969

Cerebrospinal Fluid Dynamics Studied With Radionuclides

Human serum albumin tagged with iodine-131 is currently being employed to evaluate the cerebrospinal fluid spaces in a number of neuro-

pathologic conditions including communicating hydrocephalus, CSF rhinorrhea and otorrhea, arachnoidal cysts and other obstructing lesions resulting from a variety of causes. The albumin tracer is injected in radionuclide cisternography via a lumbar puncture and the upward flow and symmetrical distribution of the tracer over the brain surface is observed.

Evaluation of the interventricular flow of the CSF is accomplished by injecting the radioactive tracer directly into the ventricular system in a manner similar to that used in air ventriculography. In addition to demonstrating altered flow caused by obstructing lesions, radionuclide ventriculography is ideally suited for evaluating the patency of neurosurgical shunts, particularly in children, where shunt revision is occasionally necessary. Shunt patency can also be examined by injecting the tracer directly into the subcutaneous pump of many of the currently used ventriculo-vascular shunts.

WILLIAM L. ASHBURN, M.D.

REFERENCES

- Ashburn WL, DiChiro G: *Radioisotope cisternography and ventriculography In Radionuclide applications in neurology and neurosurgery*. Edited by Wang and Paoletti. Springfield, Illinois, CC Thomas, 1970
- Tator CH, Fleming JFR, Sheppard RH, et al: Radioisotope test for communicating hydrocephalus. *J Neurosurg* 28:327-340, 1968

Isotope Lung Scanning In Pediatric Respiratory Disease

Although lung scanning with isotopes is a well established diagnostic procedure, its use has been greater in adults than in children, and particularly helpful in evaluation of pulmonary infarcts. Pendarvis and Swischuk presented a well-structured multi-method analysis of respiratory diseases in childhood utilizing the short half-life isotopes of indium, 113 iron hydroxide and the gamma camera scanning equipment. In addition to the isotope scanning, conventional radiography, arteriography and bronchography were used in each case.

Among conditions studied and reported were primary pulmonary artery anomalies, chronic respiratory disease such as tuberculosis, hilar and mediastinal masses, bronchiectasis, cystic fibrosis and congenital bullae.

Although isotope scanning is not a primary absolute diagnostic procedure, it is emerging in pediatrics as an increasingly useful correlative